

Antibiotic Susceptibility of Common Organisms

2022 STHS Heart - All Units

Numbers indicate percent susceptible; only the first isolate per patient per period is included

		# of isolates	Beta-Lactams							FQ	AGs	Miscellaneous				Urine Isolates Only									
			Oxacillin	Ampicillin ¹	Ampicillin/Sulbactam	Piperacillin/Tazobactam	Cefazolin	Ceftriaxone	Cefepime	Meropenem	Ciprofloxacin	Amikacin	Gentamicin	Tobramycin	Clindamycin	Tetracycline ³	TMP/SMX (Bactrim)	Vancomycin	Lincosolid	# of isolates	1st Gen. Ceph's (oral) ⁴	Nitrofurantoin ⁵	Ciprofloxacin	Ampicillin	TMP/SMX (Bactrim)
Gram Negative	Enterobacter cloacae ⁶	19	-	-	-	53	0	-	95	89	84	100	89	89	-	-	79	-	-	-	-	-	-	-	-
	Escherichia coli	290	-	31	48	92	70	74	74	100	54	100	79	74	-	-	58	-	-	229	72	96	55	33	57
	Klebsiella pneumoniae	109	-	0	56	84	65	67	68	96	78	98	83	73	-	-	62	-	-	72	75	47	89	1	67
	Proteus mirabilis	59	-	49	64	98	85	98	100	98	98	100	83	83	-	-	80	-	-	31	87	0	97	58	87
	Pseudomonas aeruginosa	77	-	-	-	83	-	-	88	86	90	97	97	100	-	-	-	-	-	8 ^Y	-	-	88	-	-
Gram Positive	Staphylococcus aureus ^{7,8}	90	62	-	-	-	-	-	-	-	-	-	94	-	73	83	86	100	100	-	-	-	-	-	-
	Staphylococcus epidermidis	35	40	-	-	-	-	-	-	-	-	-	83	-	54	77	80	97	100	-	-	-	-	-	-
	Enterococcus faecalis	80	-	100	-	-	-	-	-	-	-	-	-	-	28	-	99	94	-	-	100	76	100	-	-

Resistance: 38% MRSA, 5% VRE, 26% ESBL

¹ Ampicillin susceptibility for Enterococcus indicates susceptibility to amoxicillin, amoxicillin-clavulanate, ampicillin-sulbactam, and piperacillin-tazobactam

² Aztreonam use is reserved for severe penicillin allergy (e.g. anaphylaxis); caution: may cross-react with ceftazidime allergy

³ Tetracycline susceptibility indicates susceptibility to doxycycline and minocycline; some organisms resistant to tetracycline may be susceptible to doxycycline or minocycline

⁴ When used for therapy of uncomplicated UTIs due to E. coli, K. pneumoniae, and P. mirabilis, cefazolin susceptibility indicates susceptibility to the oral agents cefaclor, cefdinir, cefpodoxime, cefprozil, cefuroxime axetil, and cephalexin

⁵ Nitrofurantoin is indicated for use in cystitis only; it should not be used for systemic infections, including pyelonephritis

⁶ Use of 3rd generation cephalosporins is not recommended, these organisms frequently become resistant during therapy

⁷ Oxacillin and cefazolin are considered first-line antibiotic therapy for MSSA

⁸ For staphylococci and enterococci that test susceptible, aminoglycosides must be used in combination with beta-lactams or vancomycin for synergy

^Y Susceptibility rates determined on small sample size (<30) and may be statistically unreliable; interpret with caution

